

Sample Brief

Marine microbial ecology and bioreactors

By Roger Reemer, Wageningen University and Research Centre (WUR)

The Rio+20 outcome document (§158) expressed a general commitment to protect and restore marine ecosystems: “...We therefore commit to protect, and restore, the health, productivity and resilience of oceans and marine ecosystems...” However, no reference was made to the marine microbial ecology or the potentials of bioreactors. The following science digest provides an overview of scientific findings to support an informed discussion among decision-makers in the follow-up to Rio+20.

Introduction

The oil spill in the Gulf of Mexico in 2010, where the equivalent of 4.9 million barrels of oil was released into the environment, attracted a lot of attention among policymakers and civil society. However, the impacts on the ecosystem turned out to be less catastrophic than expected due to marine microbes digesting the oil spill, even though there are signs of permanent damage (Walsh 2011). This is a clear example of the adaptive properties of bacteria and the wide range of compounds they can feed on (Löffler & Edwards 2006). Microbes also have a very diverse range of substances they can produce and are at the base of healthy, stable, ecosystems all over the world. Marine microbial ecology, while still a relatively new field of research, is rapidly uncovering the importance of microbial life in nutrient availability in ecosystems. This is no different for marine environments in which, for example, processes such as nitrification (Baker et al. 2012) and CO₂ fixation (Emerson et al. 2013) are mainly regulated by microorganisms. Considering that 90 per cent of the ocean’s biomass is estimated to be microbial life (Ausubel et al. 2010), it is no surprise that microbes play a crucial role in ecosystems. However, because this field has only become an important topic in the last 10 years or so, in-depth research is still lacking and only general findings exist in terms of marine biodiversity (Millennium Ecosystem Assessment 2005) or in relation to human health (Dunn et al. 2013).

Marine microbial ecology and bioreactors: facts and figures

- Over 90% of marine biomass consists of microbial life.
- An estimated 50-80% of all biomass is found under the ocean surface.
- Microbial phytoplankton makes up the basis of the marine food chain and is responsible for producing 50% of the world’s oxygen.
- Microorganisms are also the main drivers behind nutrient availability in marine ecosystems.
- However, microbial community dynamics are still poorly understood.

Source: Authors’ compilation.

Scientific debate

Within the scientific community there is still debate on the determination of microbial species, because there is little genetic material to work with (Caron et al. 2009). In addition, one of the difficulties is determining marine microbial biodiversity, as the sheer number of species is very high and data from different areas are not properly integrated (Duffy et al. 2013). There are also the technical difficulties of measuring a large area like the ocean, which naturally comes forth from the trade-off of either covering a large area or getting a detailed description. However, it is becoming clearer that high microbial biodiversity is not necessarily the main reason for a healthy ecosystem. Rather, the composition of a certain microbial community may be a stronger indicator than biodiversity (Amaral-Zettler et al. 2010), and microbial community dynamics change when influenced by other factors, such as an increase in CO₂ levels (Brussaard et al. 2013). As such, a change in a microbial community

might have a big effect on the ecosystem's nutrient cycles and thus on the organisms living in it.

There have been suggestions of connecting the current assessments of marine biodiversity into a single global assessment on marine biodiversity. Such a systems approach is needed to establish the importance and function of microbial communities in ecosystems. This knowledge could be applied to increasing ecosystem resilience or assisting in ecosystem restoration, such as in the case of oil spills. This would not only help in improving the current assessments on marine life, but also in linking the data on marine ecosystem health at different levels *"from microbes to whales, [...] to entire ecosystems"*.

Where, on one hand microbial communities are to be studied and researched in order to determine their interactions with the ecosystem, on the other there is an ever-growing interest among the scientific community in bioreactors. A bioreactor is a machine that optimizes a natural environment for the growth of specific microbial species and communities. The marine bioreactors focus on microbial life that needs such specific living conditions (high salt concentrations, high pressure, etc.), that they cannot be cultured in a laboratory (Zhang et al. 2013). By positioning a bioreactor off the coast on the sea floor, the bioreactor's microbial life is able to thrive under its natural conditions. These bioreactors could even lead to a system in which the ocean is used in the sustainable production of medicine or other chemical substances, clean energy or even food. Moreover, increased use of bioreactors could lead to production of energy or biological compounds in a sustainable manner without damaging the ecosystem where the bioreactor is positioned.

In short, more efficient research into microbial communities and their interactions with the environment can be attained through biodiversity assessments. This could lead to better use of bioreactor technology. Finally, a better understanding of microbial ecology can help us in many fields, from ecosystem resilience and restoration to even a higher yield in seafood production.

Food for thought on marine microbial ecology and bioreactors

- Microbial communities, while not yet completely understood, are at the base of a healthy ecosystem.
- A global biodiversity assessment network helps in understanding the dynamics in microbial communities.
- A good understanding of microbial community dynamics can lead to new forms of ecosystem restoration and increasing resilience.
- Bioreactors combine the "special talents" of certain microbial species with the native marine environment to produce chemicals and energy without damaging the ecosystem.
- Stimulation of use of bioreactors can lead to an enhanced use of the ocean for sustainable production.

Source: Authors' compilation.

Issues for further consideration

The following issues were suggested by the team of young researchers for consideration by policymakers:

- Establish a global assessment on marine biodiversity, with special attention to microbial biodiversity.
- Promote research on the application of bioreactors in marine environments.
- Improve understanding and functioning of marine microbial communities.

References

- Amaral-Zettler, L., Felipe Artigas, L., Baross, J., Bharathi, L., Boetius, A., Chandramohan, D., Herndl, G., Kogure, K., Neal, P., Pedrós-Alió, C., Ramette, A., Schouten, S., Stal, S., Thessen, A., Leeuw, J., Sogin, M., (2010). A Global Census of Marine Microbes. Wiley-Blackwell, *Life in the World's Oceans: Diversity, Distribution and Abundance*. From <http://comlmaps.org/mcintyre/ch12>
- Ausubel, J. H., Trew Crist, D. & Waggoner, P. E. (2010). FIRST CENSUS OF MARINE LIFE 2010 HIGHLIGHTS OF A DECADE OF DISCOVERY. *Census of Marine Life*. From <http://www.coml.org/pressreleases/census2010/PDF/Highlights-2010-Report-Low-Res.pdf>
- Baker, B. J., Lesniewski, R. A., & Dick, G. J. (2012). Genome-enabled transcriptomics reveals archaeal populations that drive nitrification in a deep-sea hydrothermal plume. *The ISME journal*, 6(12), 2269–79. doi:10.1038/ismej.2012.64
- Brussaard, C. P. D., Noordeloos, A. A. M., Witte, H., Collenieur, M. C. J., Schulz, K., Ludwig, A., & Riebesell, U. (2013). Arctic microbial community dynamics influenced by elevated CO₂ levels. *Biogeosciences*, 10(2), 719–731. doi:10.5194/bg-10-719-2013
- Caron, D. A., Countway, P. D., Savai, P., Gast, R. J., Schnetzer, A., Moorthi, S. D., Jones, A. C. (2009). Defining DNA-based operational taxonomic units for microbial-eukaryote ecology. *Applied and environmental microbiology*, 75(18), 5797–808. doi:10.1128/AEM.00298-09
- Duffy, J. E., Amaral-Zettler, L. A., Fautin, D. G., Paulay, G., Tatiana, A., Sosik, H. M., Ryneerson, T. A. (2013). Envisioning a Marine Biodiversity Observation Network. *BioScience*, 63(5), 350–361. doi:10.1525/bio.2013.63.5.8
- Dunn, G., Harris, L., Cook, C., & Prystajek, N. (2013). A comparative analysis of current microbial water quality risk assessment and management practices in British Columbia and Ontario, Canada. *The Science of the total environment*, 468-469C, 544–552. doi:10.1016/j.scitotenv.2013.08.004
- Emerson, D., Field, E. K., Chertkov, O., Davenport, K. W., Goodwin, L., Munk, C., Woyke, T. (2013). Comparative genomics of freshwater Fe-oxidizing bacteria: implications for physiology, ecology, and systematics. *Frontiers in microbiology*, 4(September), 254. doi:10.3389/fmicb.2013.00254
- Löffler, F. E., & Edwards, E. A. (2006). Harnessing microbial activities for environmental cleanup. *Current opinion in biotechnology*, 17(3), 274–84. doi:10.1016/j.copbio.2006.05.001
- Millennium Ecosystem Assessment, 2005. *Ecosystems and Human Well-being: Biodiversity Synthesis*. World Resources Institute, Washington, DC.
- Walsh, B. (2011). The BP Oil Spill, One Year Later: How Healthy Is the Gulf Now? From <http://content.time.com/time/health/article/0,8599,2066031,00.html>
- Zhang, Y., Arends, J. B. A, Van de Wiele, T., & Boon, N. (2011). Bioreactor technology in marine microbiology: from design to future application. *Biotechnology advances*, 29(3), 312–21. doi:10.1016/j.biotechadv.2011.01.004